

- Crump BC**, Adams HE, Hobbie JE, Kling GW. 2007. Biogeography of freshwater bacterioplankton in lakes and streams of an Arctic tundra catchment. [Ecology 88: 1365–1378](#)
- Judd KE, **Crump BC**, Kling GW (2007) Bacterial responses in activity and community composition to photo-oxidation of dissolved organic matter from soil and surface waters. [Aquatic Sciences 69:96-107](#)
- Kan J, **Crump BC**, Wang K, Chen F (2006) Bacterioplankton community in Chesapeake Bay: Predictable or random assemblages. [Limnol and Oceanogr 51:2157-2169](#)
- Judd KE, **Crump BC**, Kling GW (2006) Variation in dissolved organic matter controls bacterial production and community composition. [Ecology 87:2068-2079](#)
- Crump BC**, Hobbie JE (2005) Synchrony and seasonality in bacterioplankton diversity of two temperate rivers. [Limnol and Oceanogr 50:1718-1729](#)
- Bahr M, **Crump BC**, Hobbie JE, Teske A. (2005) Molecular characterization of sulfate-reducing prokaryotes during an annual cycle in a New England salt marsh. [Environmental Microbiology 7\(8\):1175-1185](#)
- Crump BC**, Hopkinson CS, Sogin ML, Hobbie JE (2004) Microbial biogeography along an estuarine salinity gradient: the combined influences of bacterial growth and residence time. [Appl Environ Microbiol 70:1494-1505](#)
- Klepac-Ceraj V, Bahr M, **Crump BC**, Teske AP, Hobbie JE, Polz MF (2004) High overall diversity and dominance of microdiverse relationships in salt marsh sulphate-reducing bacteria. [Environmental Microbiology 6:686-698](#)
- Zwart G, van Hannen EJ, Kamst-van Agterveld MP, van der Gucht K, Lindström ES, Van Wichelen J, Lauridsen T, **Crump BC**, Han SK, Declerck S (2003) Rapid screening for freshwater bacterial groups using reverse line blot hybridization. [Appl Environ Microbiol 69: 5875-5883](#)
- Crump BC**, Bahr M, Kling GW, Hobbie JE (2003) Bacterioplankton community shifts in an arctic lake correlate with seasonal changes in organic matter source. [Appl Environ Microbiol 69:2253-2268](#)
- Levine UY, **Crump BC** (2002) Bacterioplankton community composition in flowing waters of the Ipswich River watershed. [Biol Bull 203:251-252](#)
- Zwart G, **Crump BC**, Kamst-van Agterveld MP, Hagen F, Han SK (2002) Typical freshwater bacteria: an analysis of available 16S rRNA gene sequences from plankton of freshwater lakes and rivers. [Aquat Microb Ecol 28:141-155](#)
- Crump BC**, Baross JA (2000) Characterization of the bacterially-active particle fraction in the Columbia River estuary. [Mar Ecol Prog Ser 206:13-22](#)
- Crump BC**, Baross JA (2000) Archaeaplankton in the Columbia River, its estuary, and the adjacent coastal ocean, USA. [FEMS Microbiol Ecol 31:231-239](#)
- Crump BC**, Armbrust EV, Baross JA (1999) Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary, and the adjacent coastal ocean. [Appl Environ Microbiol 65:3192-3204](#)
- Crump BC**, Baross JA, Simenstad CA (1998) Dominance of particle-attached bacteria in the Columbia River estuary, USA. [Aquatic Microbial Ecology 14:7-18](#)
- Prahl FG, Small LF, Sullivan BA, Cordell J, Simenstad CA, **Crump BC** and Baross JA (1998) Biogeochemical gradients in the lower Columbia River. [Hydrobiologia 361:37-52](#)

- Hopkinson C, Buffam I, Hobbie J, Vallino J, Perdue M, Eversmeyer B, Prahl F, Covert J, Hodson R, Moran MA, Smith E, Baross J, **Crump B**, Findlay S, Foreman K (1998) Terrestrial inputs of organic matter to coastal ecosystems: an intercomparison of chemical characteristics and bioavailability. [Biogeochemistry 43:211-234](#)
- Crump BC**, Baross JA (1996) Particle-attached bacteria and heterotrophic plankton associated with the Columbia River estuarine turbidity maxima. [Marine Ecology Progress Series 138:265-273](#)
- Baross JA, **Crump BC**, Simenstad CA (1994) Elevated 'microbial loop' activities in the Columbia River estuarine turbidity maximum. In: Dyer KR, Orth RJ (eds.) Changes in fluxes in estuaries: Implications from science to management (ECSA/ERF symposium, Plymouth, September 1992). Olsen & Olsen, Fredenborg, Denmark, p 459-464
- Pledger RJ, **Crump BC**, Baross JA (1994) A barophilic response by two hyperthermophilic, hydrothermal vent *Archaea*: An upward shift in the optimal temperature and acceleration of growth rate at supra-optimal temperatures by elevated pressure. [FEMS Microbiology Ecology 14:233-242](#)
- Erauso G, Reysenbach A-L, Godfroy A, Meunier J-R, **Crump B**, Partensky F, Baross JA, Marteinsson V, Barbier G, Pace NR, Prieur D (1993) *Pyrococcus abyssi* sp. nov., a new hyperthermophilic archaeon isolated from a deep-sea hydrothermal vent. Archives of Microbiology 160:338-349
- Baross JA, Holden JF, Pledger RJ, **Crump BC**, Summit M (1993) Submarine vent archaea: ecology and strategies for growth above 90°C. Proceedings of the International Summer Seminar on Deep-sea microorganisms. Japan Marine Science and Technology Center, Deep Star Group, JAMSTEC